



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## A SHORT DEMONSTRATION OF THE EXPONENTIAL THEOREM.

By MR. J. M. SCHAEBERLE, Ann Arbor, Mich.

To demonstrate the Exponential Theorem without the aid of the calculus, various methods have been devised by mathematicians; but all of the methods known to the writer are much longer than the one here given, which, for brevity and directness of proof, leaves but little to be desired.

Assuming that the equation

$$a^x = c_0 + c_1x + c_2x^2 + c_3x^3 + c_4x^4 + \dots \quad (1)$$

is true for all values of  $x$ , then  $c_0$  must equal unity. Squaring (1) we have

$$a^{2x} = 1 + 2c_1x + (2c_2 + c_1^2)x^2 + (2c_3 + 2c_2c_1)x^3 + (2c_4 + 2c_3c_1 + c_2^2)x^4 + \dots, \quad (2)$$

while, if in equation (1) we substitute  $2x$  in place of  $x$ , we have

$$a^{2x} = 1 + 2c_1x + 4c_2x^2 + 8c_3x^3 + 16c_4x^4 + \dots \quad (3)$$

Equating the coefficients of the like powers of  $x$  in (2) and (3), we at once obtain

$$c_1 = c_1; \quad c_2 = \frac{c_1^2}{2!}; \quad c_3 = \frac{c_1^3}{3!}; \quad c_4 = \frac{c_1^4}{4!}; \text{ etc.};$$

and if the succeeding values of  $c$  follow the same law,

$$c_{n-1} = \frac{c_1^{n-1}}{(n-1)!}, \quad c_n = \frac{c_1^n}{n!},$$

which is readily proved by equating the coefficients of the  $n$ th power of  $x$  in the second members of (2) and (3). Thus

$$2c_n + 2c_{n-1}c_1 + 2c_{n-2}c_2 + 2c_{n-3}c_3 + \dots + 2c_{\frac{1}{2}(n+1)}c_{\frac{1}{2}(n-1)} \text{ (or } c_{\frac{1}{2}n}^2) = 2^n c_n, \quad (4)$$

which, if we substitute the assumed values of  $c_n, c_{n-1}, c_{n-2}$ , etc., becomes

$$\frac{c_1^n}{n!} [2 + 2\binom{n}{1} + 2\binom{n}{2} + 2\binom{n}{3} + \dots + 2\binom{n}{\frac{1}{2}(n-1)} \text{ or } \binom{n}{\frac{1}{2}n}] = 2^n \frac{c_1^n}{n!}. \quad (5)$$

When  $n$  is even, the last term within the [ ] is  $\binom{n}{\frac{1}{2}n}$ , and when  $n$  is odd, the last term is  $2\binom{n}{\frac{1}{2}(n-1)}$ . Now, as the value of the expression within the [ ] in equation (5) is always equal to  $2^n$ , it follows that, in general,  $c_n$  must always equal  $\frac{c_1^n}{n!}$ ; and hence

$$a^x = 1 + c_1x + \frac{c_1^2x^2}{2!} + \frac{c_1^3x^3}{3!} + \frac{c_1^4x^4}{4!} + \dots + \frac{c_1^nx^n}{n!} + \dots$$